

Table 1: AIC model summary. Δt is the amount of postbounce time covered in each model, $h_{+,max}$ is the maximum axisymmetric gravitational wave strain amplitude scaled to 10 kpc distance, $h_{char,max}$ is the maximum of the characteristic strain spectrum (Flanagan & Hughes, PRD 57, 4535 (1998)) and f_{peak} is the frequency at which $h_{char,max}$ is located. E_{GW} is the total energy radiated in gravitational waves. The gravitational wave emission data for fluid matter and neutrino sources are given separately. The GW emission from anisotropic neutrino emission was computed in a postprocessing step and only coarse time resolution was available. The results of the fourier analysis of these data should be considered with this in mind.

Model	Δt (ms)	Fluid				Neutrinos			
		$ h_{+,max} $ at 10 kpc (10^{-21})	$h_{char,max}$ at 10 kpc (10^{-21})	f_{peak} (Hz)	E_{GW} ($10^{-10} M_{\odot} c^2$)	$ h_{+,max} $ at 10 kpc (10^{-21})	$h_{char,max}$ (10^{-21})	f_{peak} (Hz)	E_{GW} ($10^{-10} M_{\odot} c^2$)
1.46 M_{\odot}	510	0.55	1.45	431	2.17	4.59	1.92	$\sim 0-5$	0.01
1.92 M_{\odot}	776	3.62	2.80	166	13.16	19.88	5.83	$\sim 0-5$	0.21